

Minimal universal crossed extensions of superperfect groups

Nika Logua

e-mail: nika.logua754@ens.tsu.edu.ge

Department of Mathematics, Faculty of Exact and Natural Sciences,
Ivane Javakhishvili Tbilisi State University,
University St.2, Tbilisi 0177, Georgia

One of the basic facts in homological algebra is that every perfect group G , i. e. such that its first homology group $H_1(G)$ is trivial, possesses a central extension of G by $H_2(G)$.

Another basic fact is that the third cohomology group $H^3(G; M)$ classifies crossed extensions

$$0 \rightarrow M \rightarrow X \rightarrow Y \rightarrow G \rightarrow 1$$

where $\partial: X \rightarrow Y$ is a crossed module, with the resulting action of G on M again given by the G -module structure on M . If this action is trivial and G is superperfect, that is, both $H_1(G)$ and $H_2(G)$ are trivial, then $H^3(G; M)$ becomes isomorphic to $\text{Hom}(H_3(G), M)$, so that there exists a universal crossed extension of G by $H_3(G)$.

However, unlike the universal central extension, the universal crossed extension determined by $H_3(G)$ is not unique anymore. There is an initial representative for it, with Y a free group and X freely (in a certain sense) built from Y and the universal cocycle on $H_3(G)$, but there are many others. In particular, it follows from the classical work of Holt that if G is finite, then there is a representative with also Y finite and X finite abelian. In fact, X can be taken the induced G -module $H_3(G)[G]$.

A natural question then arises whether there exists a smallest representative, i. e. one which is a quotient of any other representative. Indeed, we will see that for any Y -subgroup $Z \subset X$ such that $M \cap Z = 0$ one can further reduce to $X/Z \rightarrow Y/\partial(Z)$, so that if a smallest representative exists, there exists a largest Z with $M \cap Z = 0$.

Our main result is that such largest Z need not exist. Namely, we show that for G equal to the Mathieu group M_{11} there exist two distinct maximal submodules $Z \subset H_3(G)[G]$ with $H_3(G) \cap Z = 0$. As a corollary we deduce that in general there is no smallest in the above sense representative for the universal crossed extension.